## Title:Frequency and characteristics of anthracycline-induced cardiotoxicities in a developing

## country in South East Asia.

**Authors:** Siew Lian Leong<sup>1,2</sup>\*, Kian Meng Chang<sup>3</sup>, Oteh Maskon<sup>4</sup>, Zhen Ying Kong<sup>5</sup>, Habiba Nazeera Begum Kamarul Jaman<sup>3</sup>, S Fadilah Abdul Wahid<sup>6</sup>, Su Shan Kong<sup>7</sup>, Ly Sia Loong<sup>8</sup>, Shaun Wen Huey Lee<sup>1,9,10</sup>

<sup>1</sup>School of Pharmacy, Monash University Malaysia, Selangor, Malaysia.

<sup>2</sup>Faculty of Pharmacy, University of Cyberjaya, Selangor, Malaysia.

<sup>3</sup>Department of Haematology, Hospital Ampang, Selangor, Malaysia.

<sup>4</sup>Cardiology Unit, Department of Medicine, Hospital CanselorTuanku Muhriz/ Pusat Perubatan UKM, Kuala Lumpur Malaysia.

<sup>5</sup>Pharmacy Department, University Malaya Medical Centre, Kuala Lumpur, Malaysia.

<sup>6</sup>Pusat Terapi Sel, Hospital CanselorTuanku Muhriz / UniversitiKebangsaan Malaysia Medical Center, Kuala Lumpur Malaysia.

<sup>7</sup>Department of Pharmacy, Hospital Ampang, Selangor, Malaysia.

<sup>8</sup>Department of Pharmacy, Hospital CanselorTuanku Muhriz/ Pusat Perubatan UKM, Kuala Lumpur Malaysia.

<sup>9</sup>Asian Centre for Evidence Synthesis in Population, Implementation and Clinical Outcomes (PICO), Health and Well-being Cluster, Global Asia in the 21st Century (GA21) Platform, Monash University Malaysia, Selangor, Malaysia.

<sup>10</sup>School of Pharmacy, Taylor's University Lakeside Campus, Selangor, Malaysia.

## \*Corresponding author:

Siew Lian Leong, Faculty of Pharmacy, University of Cyberjaya Persiaran Bestari, Cyber 11, 63000 Cyberjaya, Selangor, Malaysia. siewlian@cyberjaya.edu.my Tel: +601-2881 7198

## Abstract:

Background: The prevalence of anthracyclines-induced cardiotoxicity differs between regions and

countries, but little is known about the incidence in low-middle income countries including Malaysia.

Objective: To determine the incidence and characteristics of anthracycline-induced cardiotoxicity

among cancer patients in Malaysia.

Setting: A retrospective-observational study in three tertiary hospitals in Klang Valley, Malaysia.